this month in diseases

## By A.R. Chase and Margery Daughtrey

T	Disease	Pathogen	Stage of crop	Best cultural controls	Best product controls, based mainly on Chase Research trials	LX
	Alternaria leaf spot	<i>Alternaria</i> spp.	Finishing and landscape	Avoid overhead irrigation.	Daconil Ultrex, strobilurins (like Compass O, Heritage and Insignia)	In this issue, we present a summary of diseases that occur in petunias. Remember: Product controls are based on research trials and do not reflect an endorsement of any sort.
	Black root rot	Thielaviopsis basicola	From plugs to finish	Maintain potting medium pH.	Thiophanate methyl products (like 3336 and OHP-6672) and Veranda O	
	Botrytis blight and stem rot	Botrytis cinerea	Anytime from plug production to finishing	Irrigation management; don't grow plants under hanging baskets that drop petals.	Daconil Ultrex (before flowering starts), Chipco 26019, Decree, Medallion and Pageant	
	Mosaic Virus	Tobacco Mosaic Virus	Anytime from plug to finishing	Use virus-indexed cuttings; discard all infected plants.	Clean benches etc. between crops with nonfat dry milk (20 percent) or bleach (10 percent).	
	Phytophthora aerial blight	Phytophthora nicotianae (=parasitica)	Finishing and landscape	Do not plant ground beds consecutively with the same crop.	Subdue MAXX (resistance is possible), Terrazole (or Truban), Aliette (or other phosphonates) and Stature	
	Powdery mildew	<i>Podosphaera xanthii</i> and <i>Oidium</i> <i>longipes</i>	Sometimes found in spring or in landscape at end of season	Scout regularly and maintain spacing and air movement.	Sterol inhibitors (like Eagle) and strobilurins (like Compass O, Cygnus and Insignia)	- A.R. Chase is president and pathologist of Chase Horticultural Research, Inc. and can be reached at archase@ chaseresearch.net. Margery L. Daughtrey is senior extension associate at Cornell University's Long Island Horticultural Research & Extension Center and can be reached at mld9@ cornell.edu.
	Pythium damping-off and root rot	Pythium irregulare	Plug production mainly	Use well-draining potting media; manage water carefully; use new potting media and pots.	Subdue MAXX (resistance is possible), Terrazole (or Truban), FenStop (use lower labeled rates) and Segway	
	Rhizoctonia stem rot	Rhizoctonia solani	All stages of production	Use new rooting media and flats.	Medallion, thiophanate methyl (like OHP-6672 or 3336), and strobilurin (like Heritage) sprench	
	Rhizopus blight	Rhizopus stolonifer	Finishing under extreme heat	Maintain spacing and scout; reduce stress during extreme heat.	Treatments with Medallion have been effective on other crops.	
TROCK	Sclerotinia blight	Sclerotinia sclerotiorum	Finishing in spring, mainly	Maintain spacing for good air movement.	Palladium and Pageant	